



Research Article

Prevalence of Abnormal Body Posture Among Chinese Children and Adolescents: A Large Population (158.584) -Based Cross-Sectional Study

 Lei Yang,¹  Xinhai Lu,¹⁻³  Bin Yan,¹⁻³  Yeen Huang¹⁻³

¹Department of Spine Surgery, The First Affiliated Hospital of Shenzhen University, Shenzhen, China

²Department of Spine Surgery, The Shenzhen Second People's Hospital, Shenzhen, China

³Department of Spine Surgery, Shenzhen Youth Spine Health Center, Shenzhen, China

Abstract

Objectives: Poor body posture in adulthood is often formed from the childhood, and individuals with severe abnormal body posture may associated with the progress to scoliosis. However, there is still a lack of epidemiological evidence in the prevalence of adolescent abnormal body posture compared to scoliosis.

Methods: We sample a large representative dataset comprising data collected from 1th to 12th graders using a multistage, stratified-cluster, random-sampling method. Our sample consisted of 158.584 children and adolescents who were screened by visual inspection of clinical signs, the forward-bending test, and the measurement of angle of trunk rotation (ATR).

Results: The overall prevalence of abnormal body posture in Chinese children and adolescents was 65.3%, around 3.7% of children and adolescents were referred for radiography. According to gender and age, girls had a higher prevalence of abnormal body posture than boys (76.0% vs. 56.6%, $p < 0.001$), students aged 10-15 and >15 years old had a higher prevalence of abnormal body posture than students <10 years old (64.8% and 71.1% vs. 41.3%, $p < 0.001$).

Conclusion: Our findings showed that Chinese children and adolescents have a significantly higher prevalence of abnormal body posture, girls and older students may be an especially high-risk group.

Keywords: abnormal body posture, adolescent, children, prevalence, school screening program

Cite This Article: Yang L, Lu X, Yan B, Huang Y. Prevalence of Abnormal Body Posture Among Chinese Children and Adolescents: A Large Population (158.584) -Based Cross-Sectional Study. EJMO 2020;4(1):71–78.

Abnormal body posture refers to an abnormal body state in which the body cannot maintain a stable state and the normal function of tissues and organs in an upright state.^[1] Previous studies have shown that the poor body posture in adulthood is often formed from the childhood.^[2] Children and adolescents with severe abnormal body posture may associated with the progress to adolescent idiopathic scoliosis (AIS).^[3,4] Understanding the current prevalence of abnormal body posture of children and adolescents will help to identify high-risk population and formulate targeted interventions.

Previous studies showed that 34%-50% of children and

adolescents have different degrees of abnormal body posture,^[5,6] less than a fifth of children and adolescents have a normal body posture,^[7] most of the children have certain abnormal posture problems (e.g. shoulder asymmetry, kyphosis, or scapula tilt). Furthermore, some researchers suggested that compared with children of the same age who have not received school education, the students who have received school education have more serious body posture problems.^[8,9]

To the best of our knowledge, previous evidence mainly come from the western or developed countries. In China, only one school screening program conducted in Beijing

Address for correspondence: Bin Yan, M.M. Department of Spine Surgery, The First Affiliated Hospital of Shenzhen University, Shenzhen, China

Phone: +86 0755-83243394 **E-mail:** huangyeenzs@163.com

Submitted Date: December 12, 2019 **Accepted Date:** January 31, 2020 **Available Online Date:** March 12, 2020

©Copyright 2020 by Eurasian Journal of Medicine and Oncology - Available online at www.ejmo.org

OPEN ACCESS This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.



showed that nearly 80% of children and adolescents were reported to have at least one sign of abnormal body postures.^[10] However, there is still a lack of large-scale epidemiological evidence in the prevalence of adolescent abnormal body posture.

Therefore, we conducted this large-scale cross-sectional study in south China to estimate the prevalence of abnormal body posture among children and adolescents, to describe the epidemiological findings according to the demographic characteristics, to provide data support for finding high-risk population and targeted interventions.

Methods

Study Design and Participants

This is a school-based cross-sectional study in children and adolescent students from the Shenzhen city in south China. All students were selected by a 3-stage, stratified-cluster, random-sampling method. In stage 1, we divided Shenzhen city into three economic stratifications (high-level, middle-level, and low-level) by per capita GDP (gross domestic product), and then selected two districts from each stratification by simple randomization using random number generator. In stage 2, schools in each district were divided into three categories: primary schools (i.e., grades 1-6), junior high schools (i.e., grades 7-9), and senior high schools (i.e., grades 10-12). Based on the proportions of these three types of schools, six primary schools, six junior high schools, and three senior high schools were randomly selected from each representative district. In stage 3, all classes from each grade within the selected schools were included in our study, and all available students in the selected classes were invited to voluntarily participate in the present study.

In total, 159,120 students were invited to participate in our scoliosis screening, 158,584 students were completed and qualified for our study.

Ethical Statement

This study was conducted in accordance with the Declaration of Helsinki, and was approved by the Shenzhen Second People's Hospital Institutional Review Board (Ethical code: SZ20130816). Written or oral informed consent was obtained from the parent or legal guardian of each participating student under 18 years old or from each participating student who at least 18 years old.

Data Collection

To protect the privacy of the students, all students in the selected classes were screened for scoliosis in a closed room or tent, and administered by research assistants without the presence of teachers or other school personnel (to avoid potential information bias). All data were collected from September to November 2019.

School Screening Program

The school screening program for AIS in Shenzhen was started in 2013 as part of the national public health project, and conducted and administered by the Shenzhen Youth Spine Health Center (SYSHC) of the Shenzhen Second People's Hospital using a national scoliosis screening standardized protocol (GB/T 16133-2014).^[11] Students in primary schools, junior high schools, and high schools were invited to participate in the screening program voluntarily. School screening was performed by an experienced team of trained rehabilitation therapists from SYSHC using the Adams forward bending test (FBT), visual inspection, and measurement of the angle of trunk rotation (ATR) using the scoliometer.^[12]

Abnormal Body Posture

Abnormal body posture was assessed by visual inspection, FBT, and ATR.^[13,14] The standard visual inspection was performed in the upright position. The examiner checked for spine alignment, shoulder asymmetry (high and low shoulder), scapula prominence (scapula tilt), hip and pelvic obliquity (pelvic tilt), back symmetry (flat back, kyphosis), lumbar curvature (lumbar concave, lumbar lordosis, and lumbar kyphosis), distance of the hands from the flanks, and length of the lower limbs.^[15] The FBT was performed with the student's feet placed together, knees straight, while bending at the hips to nearly 90 degrees with the arms freely hanging forward, palms together. Any significant clinical sign was recorded. The ATR was used to decide the frequency of assessments and when should the students be referred for radiography directly by measuring with a scoliometer, including angle of thoracic rotation, angle of lumbar rotation, and angle of thoracolumbar rotation.^[16] Students with an ATR $>5^\circ$ or with 1 or more significant clinical signs were identified as abnormal body posture.

Demographic Variables

Demographic variables included gender (boy, girl), age, ethnic group, school category, and grade. Ethnic group was assessed based on the student's self-report about their ethnic group (Han or minorities).^[17] School category included primary school (grade 1-6), junior middle school (grade 7-9), and high school (grade 10-12).

Statistical Analysis

First, descriptive analyses were conducted to describe the demographic characteristics (e.g. gender and age) of children and adolescents in south China. Second, prevalence of abnormal body posture among total population was reported, and sample was further divided into different groups to calculate gender-, age-, school-specific prevalence rates. Chi-square test (χ^2) was used to compare the

differences between groups. Bias-corrected 95% confidence intervals (CI) were estimated using 1000 bootstrap samples. All data were analyzed using SPSS 22.0 (IBM Corp, Armonk, NY, USA), a two-tailed with a P-value of less than 0.05 was considered statistically significant.

Results

Demographic Characteristics of Chinese Children and Adolescents

The sample demographic information was shown in Table 1. A total of 158,584 Chinese children and adolescents were screened, 55.3% (87754) were boys, and 44.7% were girls, yielding a male-to-female ratio of 1.2:1, and the mean (SD) age of the students was 12.8±2.0 years. Primary school, junior high school, and senior high school students account for 38.0%, 42.7%, and 19.3% respectively.

Prevalence of Abnormal Body Posture of Chinese Children and Adolescents

As shown in Table 2, the overall prevalence of abnormal body posture among Chinese children and adolescents was 65.3%

Table 1. Demographic characteristics of Chinese children and adolescents (n=158.584)

Variables	n (%)
Gender	
Boy	87754 (55.3)
Girl	70830 (44.7)
Age (Mean±SD)	12.8±2.0
Ethnic group	
Han	153886 (97.0)
Minority	4662 (3.0)
School category	
Primary school	60131 (38.0)
Junior middle school	67785 (42.7)
High school	30668 (19.3)
Grade	
Grade 1	394 (0.2)
Grade 2	401 (0.3)
Grade 3	400 (0.3)
Grade 4	461 (0.3)
Grade 5	30107 (19.0)
Grade 6	28368 (17.9)
Grade 7	24352 (15.4)
Grade 8	24008 (15.0)
Grade 9	19425 (12.2)
Grade 10	15881 (10.0)
Grade 11	14200 (9.0)
Grade 12	587 (0.4)

n: number; SD: standard deviation.

Table 2. Prevalence of abnormal body posture among Chinese children and adolescents (n=158.584)

Variables	n (%)	95% CI
Total	158584 (100.0)	
High and low shoulder		
Normal	71832 (45.3)	45.1-45.6
Left shoulder height	47935 (30.2)	30.0-30.5
Right shoulder height	38817 (24.5)	24.2-24.7
Scapula tilt		
Normal	91695 (57.8)	57.6-58.1
Tilt to the left	40007 (25.2)	25.0-25.5
Tilt to the right	26882 (17.0)	16.8-17.1
Pelvic tilt		
Normal	146165 (92.2)	92.0-92.3
Tilt to the left	6009 (3.8)	3.7-3.9
Tilt to the right	6410 (4.0)	3.9-4.2
Flat back		
Normal	157507 (99.3)	99.2-99.4
Abnormal	1077 (0.7)	0.6-0.8
Kyphosis		
Normal	154264 (97.3)	97.2-97.4
Abnormal	4320 (2.7)	2.6-2.8
Lumbar concave		
Normal	122166 (77.0)	76.8-77.2
Left concave	14754 (9.3)	9.2-9.4
Right concave	21664 (13.7)	13.5-13.8
Lumbar lordosis		
Normal	157834 (99.5)	99.4-99.6
Abnormal	750 (0.5)	0.4-0.6
Lumbar kyphosis		
Normal	158264 (99.8)	99.7-99.9
Abnormal	320 (0.2)	0.1-0.3
Angle of thoracic rotation		
Normal (ATR: 0-5°)	156239 (98.5)	98.4-98.6
Rotate to the left (ATR>5°)	696 (0.4)	0.3-0.5
Rotate to the right (ATR>5°)	1649 (1.1)	1.0-1.2
Angle of lumbar rotation		
Normal (ATR: 0-5°)	155291 (97.9)	97.8-98.0
Rotate to the left (ATR>5°)	2470 (1.6)	1.5-1.7
Rotate to the right (ATR>5°)	823 (0.5)	0.4-0.6
Angle of thoracolumbar rotation		
Normal (ATR: 0-5°)	157885 (99.6)	99.5-99.7
Rotate to the left (ATR>5°)	309 (0.2)	0.1-0.3
Rotate to the right (ATR>5°)	309 (0.2)	0.1-0.3
Abnormal body posture ^a		
Normal	55048 (34.7)	34.5-35.0
Abnormal	103536 (65.3)	65.0-65.5
Referral for radiography ^b		
No	152775 (96.3)	96.2-96.4
Yes	5809 (3.7)	3.6-3.8

n: number; CI: confidence intervals; ATR: angle of trunk rotation;

^aAbnormal body posture was defined as participant who was screened out one or more to the following abnormal physical signs: high and low shoulder, scapula tilt, pelvic tilt, flat back, kyphosis, lumbar concave, lumbar lordosis, lumbar kyphosis, angle of thoracic rotation >5°, angle of lumbar rotation >5°, angle of thoracolumbar rotation >5°; ^bReferral for radiography: an angle of trunk (thoracic, lumbar or thoracolumbar) rotation of 5° was the recommended threshold for referral to radiography.

(95% CI: 65.0%-65.5%), around 3.7% (95% CI: 3.6%-3.8%) of children and adolescents were referred for radiography. The most serious sign of abnormal body posture was high and low shoulders (left shoulder height: 30.2%, 95% CI: 30.0%-30.5%; Right shoulder height: 24.5%, 95% CI: 24.2%-24.7%) and scapula tilt (tilt to the left: 25.2%, 95% CI: 25.0%-25.5%; tilt to the right: 17.0%, 95% CI: 16.8%-17.1%).

Prevalence of Abnormal Body Posture of Chinese Children and Adolescents Stratified by Gender

According to gender (as shown in Table 3), the prevalence of abnormal body posture was significantly higher in girls than that in boys (76.0% vs. 56.6%, $\chi^2=6519.15$, $p<0.001$), girls had a 1.34 times higher prevalence of abnormal body posture than boys. Girls had a higher prevalence of referral for radiography than boys (5.2% vs. 2.4%, $\chi^2=822.31$, $p<0.001$). Besides pelvic tilt and kyphosis, the rest of the abnormal signs were higher for girls than boys.

Prevalence of Abnormal Body Posture of Chinese Children and Adolescents Stratified by Age

According to age (as shown in Table 4), the prevalence of abnormal body posture was significantly higher in students aged 10-15 and >15 years old than that in students <10 years old (64.8% and 71.1% vs. 41.3%, $\chi^2=681.18$, $p<0.001$), students aged 10-15 and >15 years old had a 1.57 and 1.72 times higher prevalence of abnormal body posture than students <10 years old, respectively. Students aged 10-15 and >15 years old had a higher prevalence of referral for radiography than students <10 years old (3.3% and 6.7% vs. 0.5%, $\chi^2=590.82$, $p<0.001$). Besides lumbar lordosis and lumbar kyphosis, the rest of the abnormal signs were higher for students aged 10-15 and >15 years old than students <10 years old.

Discussion

In this study, our large-scale cross-sectional study found that Chinese children and adolescents had serious body posture problems, around 3.7% children and adolescents were referred for radiography, girls and senior students were especially at high risk. These findings help describe the epidemiological characteristics of abnormal body posture in Chinese children and adolescents, and provide evidence for identifying high-risk groups.

Our results illustrated that the overall prevalence of abnormal body posture among children and adolescents in China was 65.3%, around 3.7% of children and adolescents were referred for radiography, which was similar to previous research findings.^[15,18] In western countries, most of the primary and secondary school students have a certain degree of abnormal body posture, only 18%-50% of chil-

dren and adolescents have a normal body posture.^[19,20] The findings of our study in the coastal areas of South China also showed the similar epidemic characteristics with the previous studies, the high and low shoulders and scapula tilt account for a high proportion, indicating that the physical posture of children and adolescents in China has become more serious, which needs to be highly concerned by education departments and related public health organizations.

Furthermore, our group comparison results showed that the prevalence of abnormal body posture in girls was much higher than that in boys, 5.2% girls were referred for radiography which was 2.2 times higher than that in boys. Our findings were consistent with Alen^[21] and Penha,^[22] but a study conducted in northern China showed that boys had a higher prevalence of abnormal body posture than that of girls.^[23] This variation in results may derive from the different sampling areas and age structures. Due to the earlier physiological development, girls are more likely to extend their neck and chest to reduce the change of appearance image.^[24,25] Besides, girls are less than boys in physical activity intensity, which tends to lead to lack of muscle strength, resulting in girls are more difficult to control body posture than boys.^[26] Therefore, based on these possible reasons, girls may be a high-risk group with abnormal body posture.

Previous studies have shown that with age, children and adolescents' body posture problems would become more serious.^[27] Our results also found that students aged 10-15 and >15 years old had more serious body posture problems than the students aged <10 years old. The reason for the rising trend of abnormal body posture rate with the aging may be related to the increase of hormone secretion and the significant changes of physical development and mental status in adolescence.^[28,29] Therefore, adolescence may be a period of high incidence of abnormal body posture.

In our large-scale school screening program conducted in south China, apart from the objective criteria (ATR of thoracic, lumbar, or thoracolumbar), a student would also be examined when there were significant signs of body appearances, including uneven shoulder height, scapular prominence, hip and pelvic obliquity, and so on.^[13,15] According to our best knowledge, this is the first population-based, cross-sectional study to report the prevalence of abnormal body posture of children and adolescents. Our findings can provide essential information to the understanding of the epidemiology of abnormal body posture, which can be an important reason for progression to adolescent idiopathic scoliosis, and to help policymakers develop appropriate program through rational planning. Our data showed that there is a high prevalence of abnormal

Table 3. Prevalence of abnormal body posture stratified by gender (n=158,584)

Variables	Gender		χ^2	P
	Boy	Girl		
Total	87754 (55.3)	70830 (44.7)		
High and low shoulder			9709.54	<0.001
Normal	48849 (55.7)	22983 (32.4)		
Left shoulder height	24001 (27.3)	23934 (33.8)		
Right shoulder height	14904 (17.0)	23913 (33.8)		
Scapula tilt			4431.13	<0.001
Normal	56530 (64.4)	35165 (49.7)		
Tilt to the left	20592 (23.5)	19415 (27.4)		
Tilt to the right	10632 (12.1)	16250 (22.9)		
Pelvic tilt			2167.83	<0.001
Normal	71393 (81.4)	63251 (89.3)		
Tilt to the left	6182 (7.0)	3719 (5.3)		
Tilt to the right	10179 (11.6)	3860 (5.4)		
Flat back			263.56	<0.001
Normal	87422 (99.6)	70085 (98.9)		
Abnormal	332 (0.4)	745 (1.1)		
Kyphosis			128.61	<0.001
Normal	84998 (96.9)	69266 (97.8)		
Abnormal	2756 (3.1)	1564 (2.2)		
Lumbar concave			2164.81	<0.001
Normal	71393 (81.4)	50773 (71.7)		
Left concave	6182 (7.0)	8572 (12.1)		
Right concave	10179 (11.6)	11485 (16.2)		
Lumbar lordosis			20.18	<0.001
Normal	87400 (99.6)	70434 (99.4)		
Abnormal	354 (0.4)	396 (0.6)		
Lumbar kyphosis			22.43	<0.001
Normal	87619 (99.8)	70645 (99.7)		
Abnormal	135 (0.2)	185 (0.3)		
Angle of thoracic rotation			383.52	<0.001
Normal (ATR: 0-5°)	86886 (99.0)	69353 (97.9)		
Rotate to the left (ATR>5°)	343 (0.4)	353 (0.5)		
Rotate to the right (ATR>5°)	525 (0.6)	1124 (1.6)		
Angle of lumbar rotation			476.74	<0.001
Normal (ATR: 0-5°)	86548 (98.6)	68743 (97.1)		
Rotate to the left (ATR>5°)	898 (1.0)	1572 (2.2)		
Rotate to the right (ATR>5°)	308 (0.4)	515 (0.7)		
Angle of thoracolumbar rotation			92.53	<0.001
Normal (ATR: 0-5°)	87492 (99.7)	70393 (99.3)		
Rotate to the left (ATR>5°)	125 (0.1)	184 (0.3)		
Rotate to the right (ATR>5°)	137 (0.2)	253 (0.4)		
Abnormal body posture ^a			6519.15	<0.001
Normal	38071 (43.4)	16977 (24.0)		
Abnormal	49683 (56.6)	53853 (76.0)		
Referral for radiography ^b			822.31	<0.001
No	85606 (97.6)	67169 (94.8)		
Yes	2148 (2.4)	3661 (5.2)		

ATR: angle of trunk rotation; ^aAbnormal body posture was defined as participant who was screened out one or more to the following abnormal physical signs: high and low shoulder, scapula tilt, pelvic tilt, flat back, kyphosis, lumbar concave, lumbar lordosis, lumbar kyphosis, angle of thoracic rotation >5°, angle of lumbar rotation >5°, angle of thoracolumbar rotation >5°; ^bReferral for radiography: an angle of trunk (thoracic, lumbar or thoracolumbar) rotation of 5° was the recommended threshold for referral to radiography.

Table 4. Prevalence of abnormal body posture stratified by age (n=158,584)

Variables	Age (year)			χ^2	P
	<10	10-15	>15		
Total	1510 (0.9)	138279 (87.2)	18795 (11.9)		
High and low shoulder				798.23	<0.001
Normal	1050 (69.5)	63376 (45.8)	7406 (39.4)		
Left shoulder height	158 (10.5)	40965 (29.6)	6812 (36.2)		
Right shoulder height	302 (20.0)	33938 (24.6)	4577 (24.4)		
Scapula tilt				203.03	<0.001
Normal	1107 (73.3)	80096 (57.9)	10492 (55.8)		
Tilt to the left	197 (13.1)	34707 (25.1)	5103 (27.2)		
Tilt to the right	206 (13.6)	23476 (17.0)	3200 (17.0)		
Pelvic tilt				101.97	<0.001
Normal	1492 (98.8)	127453 (92.2)	17220 (91.6)		
Tilt to the left	7 (0.5)	5264 (3.8)	738 (3.9)		
Tilt to the right	11 (0.7)	5562 (4.0)	837 (4.5)		
Flat back				94.57	<0.001
Normal	1506 (99.7)	137435 (99.4)	18566 (98.8)		
Abnormal	4 (0.3)	844 (0.6)	229 (1.2)		
Kyphosis				34.76	<0.001
Normal	1498 (99.2)	134410 (97.2)	18356 (97.7)		
Abnormal	12 (0.8)	3869 (2.8)	439 (2.3)		
Lumbar concave				432.13	<0.001
Normal	1365 (90.4)	107214 (77.5)	13587 (72.3)		
Left concave	62 (4.1)	12444 (9.0)	2248 (12.0)		
Right concave	83 (5.5)	18621 (13.5)	2960 (15.7)		
Lumbar lordosis				4.06	0.132
Normal	1501 (99.4)	137610 (99.5)	18723 (99.6)		
Abnormal	9 (0.6)	669 (0.5)	72 (0.4)		
Lumbar kyphosis				1.39	0.499
Normal	1509 (99.9)	137998 (99.8)	18757 (99.8)		
Abnormal	1 (0.1)	281 (0.2)	38 (0.2)		
Angle of thoracic rotation					<0.001 ^c
Normal (ATR: 0-5°)	1509 (99.9)	136425 (98.7)	18305 (97.4)		
Rotate to the left (ATR>5°)	0 (0.0)	571 (0.4)	125 (0.7)		
Rotate to the right (ATR>5°)	1 (0.1)	1283 (0.9)	365 (1.9)		
Angle of lumbar rotation				380.50	<0.001
Normal (ATR: 0-5°)	1505 (99.7)	135729 (98.2)	18057 (96.1)		
Rotate to the left (ATR>5°)	3 (0.2)	1896 (1.4)	571 (3.0)		
Rotate to the right (ATR>5°)	2 (0.1)	654 (0.4)	167 (0.9)		
Angle of thoracolumbar rotation					<0.001 ^c
Normal (ATR: 0-5°)	1508 (99.9)	137741 (99.6)	18636 (99.2)		
Rotate to the left (ATR>5°)	0 (0.0)	234 (0.2)	75 (0.4)		
Rotate to the right (ATR>5°)	2 (0.1)	304 (0.2)	84 (0.4)		
Abnormal body posture ^a				681.18	<0.001
Normal	887 (58.7)	48728 (35.2)	5433 (28.9)		
Abnormal	623 (41.3)	89551 (64.8)	13362 (71.1)		
Referral for radiography ^b				590.82	<0.001
No	1502 (99.5)	133738 (96.7)	17535 (93.3)		
Yes	8 (0.5)	4541 (3.3)	1260 (6.7)		

ATR: angle of trunk rotation; ^aAbnormal body posture was defined as participant who was screened out one or more to the following abnormal physical signs: high and low shoulder, scapula tilt, pelvic tilt, flat back, kyphosis, lumbar concave, lumbar lordosis, lumbar kyphosis, angle of thoracic rotation>5°, angle of lumbar rotation>5°, angle of thoracolumbar rotation>5°. ^bReferral for radiography: an angle of trunk (thoracic, lumbar or thoracolumbar) rotation of 5° was the recommended threshold for referral to radiography. ^cFisher's exact probability test.

body posture among Chinese children and adolescents, it is of importance to screen boys and senior students for a significantly high rate.

The present study has several limitations that are worth noting. First, due to the cross-sectional nature of the data, it is difficult to make causal inferences. Second, our study sample included only school students and did not include children and adolescents who had dropped out of school or were not present in school on the day the screening was conducted. Third, although gender and age have been reported to be important factors in abnormal body posture,^[27] other relevant influencing factors (e.g., genetics, hormone, and nutritional status)^[30–32] have not been investigated in this study.

Conclusion

Our large-scale epidemiology evidence showed that Chinese children and adolescents have a significantly higher prevalence of abnormal body posture, girls and older students may be an especially high-risk group. Appropriate prevention and intervention programs targeting for school students should be established to improve their physical health.

Disclosures

Ethics Committee Approval: The study was approved by the Shenzhen Second People's Hospital Institutional Review Board (Ethic code: SZ20130816).

Peer-review: Externally peer-reviewed.

Conflict of Interest: None declared.

Funding: This work was supported by the Scoliosis screening program for primary and secondary school students in Shenzhen (Project number: SFG [2019] No.780).

Authorship Contributions: Concept – Y.H., B.Y.; Design – Y.H.; Supervision – Y.H., B.Y.; Materials – L.Y.; Data collection &/or processing – L.Y., X.L.; Analysis and/or interpretation – L.Y.; Literature search – X.L.; Writing – L.Y., Y.H.; Critical review – Y.H.

References

- Dolphens M, Cagnie B, Coorevits P, Vanderstraeten G, Cardon G, Dhooge R, Danneels L. Sagittal standing posture and its association with spinal pain: A school-based epidemiological study of 1196 Flemish adolescents before age at peak height velocity. *Spine* 2012;37:1657. [CrossRef]
- Louw QA, Morris LD, Grimmer-Somers K. The Prevalence of low back pain in Africa: a systematic review. *BMC Musculoskelet Disord* 2007;8:105. [CrossRef]
- Nissinen M, Heliövaara M, Seitsamo J, Poussa M. Trunk asymmetry, posture, growth, and risk of scoliosis. A three-year follow-up of Finnish prepubertal school children. *Spine* 1993;18:8–13. [CrossRef]
- Nault ML, Allard P, Hinse S, Le Blanc R, Caron O, Labelle H, Sadeghi H. Relations between standing stability and body posture parameters in adolescent idiopathic scoliosis. *Spine* 2002;27:1911–7.
- Motylewski S, Zientala A, Pawlicka-Lisowska A, Poziomska-Piątkowska E. Assessment of body posture in 12- and 13-year-olds attending primary schools in Pabianice. *Pol Merkur Lekarski* 2016;39:368–71.
- Mahlknecht JF. The prevalence of postural disorders in children and adolescents: a cross sectional study. *Z Orthop Unfall* 2007;145:338–42. [CrossRef]
- Kamal, S. A. Pattern recognition using moiré fringe topography and raserstereography. *Int sym bio sec tech* 2008;1–7.
- Mahlknecht J F. The prevalence of postural disorders in children and adolescents: a cross sectional study. *Z Orthop Unfall* 2007;145:338–42. [CrossRef]
- Kasten AP, Rosa BND, Schmit EFD, Noll Matias, Candotti CT. Prevalence of postural deviations in the spine in schoolchildren: a systematic review with meta-analysis. *J Hum Grow Dev* 2017;27. [CrossRef]
- Xing Fei. Investigation and analysis on the status quo of abnormal body posture of primary and secondary school students-Taking Beijing as an example. *Gui spo sci tech* 2018;133:54–7.
- National Health and Family Planning Commission of the people's Republic of China and National Standardization Administration Commission of China. Screening of spinal curvature abnormality of children and adolescents (GB/T 16133-2014). Available online: <http://dgedu.dandong.gov.cn/docs/2019-06/20190618095730294569.pdf>. (accessed on 26 November 2019).
- Grossman D C, Curry S J, Owens D K, Barry MJ, Davidson KW, Doubeni CA, Epling JW Jr, Kemper AR, Krist AH, Kurth AE, Landefeld CS, Mangione CM, Phipps MG, Silverstein M, Simon MA, Tseng CW. Screening for Adolescent Idiopathic Scoliosis: US Preventive Services Task Force Recommendation Statement. *JAMA* 2018;319:165–72. [CrossRef]
- Hengwei F, Zifang H, Qifei W, Weiqing T, Nali D, Ping Y, Junlin Y. Prevalence of Idiopathic Scoliosis in Chinese Schoolchildren: A Large, Population-Based Study. *Spine* 2016;41:259–64.
- Fong DY, Lee CF, Cheung KM, Cheng JC, Ng BK, Lam TP, Mak KH, Yip PS, Luk KD. A meta-analysis of the clinical effectiveness of school scoliosis screening. *Spine* 2010;35:1061–71.
- Luk KD, Lee CF, Cheung KM, Cheng JC, Ng BK, Lam TP, Mak KH, Yip PS, Fong DY. Clinical effectiveness of school screening for adolescent idiopathic scoliosis: A large population-based retrospective cohort study. *Spine* 2010;35:1607–14. [CrossRef]
- Grossman TW, Mazur J M, Cummings R J. An evaluation of the Adams forward bend test and the scoliometer in a scoliosis school screening setting. *J Pediatr Orthop* 1995;15:535–8.
- Yang Z, Wang K, Li T, Sun W, Li Y, Chang YF, Dorman JS, LaPorte RE. Childhood Diabetes in China: Enormous variation by place

- and ethnic group. *Diabetes Care* 1998;21:525–9. [\[CrossRef\]](#)
18. Lee C, Fong DY, Cheung KM, Cheng JC, Ng BK, Lam TP, Mak KH, Yip PS, Luk KD. Referral criteria for school scoliosis screening: assessment and recommendations based on a large longitudinally followed cohort. *Spine* 2010;35:E1492–E1498. [\[CrossRef\]](#)
 19. Kratenová J, Zejglicová K, Malý M, Filipová V. Prevalence and risk factors of poor posture in school children in the Czech Republic. *J Sch Health* 2007;77:131–7. [\[CrossRef\]](#)
 20. Jakub P. Prevalence of postural disorders in children from Copper Basin in Poland. *Fizjoter* 2011;19:3–10. [\[CrossRef\]](#)
 21. Alen Ćirić D Č A B. Differences in posture status between boys and girls 6 to 9 years of age. *Homo Sporticus* 2015;572:12–20.
 22. Penha PJ, Penha NLJ, De Carvalho BKG, Andrade RM, Schmitt ACB, João SMA. Posture Alignment of Adolescent Idiopathic Scoliosis: Photogrammetry in Scoliosis School Screening. *J Manipulative Physiol Ther* 2017;40:441–51. [\[CrossRef\]](#)
 23. Li L. The current situation of poor body shape of primary school students in Shijiazhuang. *China school health* 2018;39:1416–8.
 24. Cash T F, Pruzinsky T. Body images: Development, deviance, and change. *Ann Plast Surg* 1992;29:367. [\[CrossRef\]](#)
 25. Feingold A, Mazzella R. Gender differences in body image are increasing. *Psy Sci* 1998;9:190–5. [\[CrossRef\]](#)
 26. Klassonheggebo L, Anderssen S A. Gender and age differences in relation to the recommendations of physical activity among Norwegian children and youth. *Scand J Med Sci Sports* 2003;13:293–8. [\[CrossRef\]](#)
 27. Ludwig O, Mazet C, Mazet D, Hammes A, Schmitt E. Age-dependency of posture parameters in children and adolescents. *J Phys Ther Sci* 2016;28:1607–10. [\[CrossRef\]](#)
 28. Richmond E, Rogol A D. Endocrine Responses to Exercise in the Developing Child and Adolescent. *Front Horm Res* 2016;58–67. [\[CrossRef\]](#)
 29. Hackney A C, Davis H C, Lane A R. Growth Hormone-Insulin-Like Growth Factor Axis, Thyroid Axis, Prolactin, and Exercise. *Front Horm Res* 2016;1–11. [\[CrossRef\]](#)
 30. Weinstein S L, Dolan L A, Cheng J C, Danielsson A, Morcuende JA. Adolescent idiopathic scoliosis. *Lancet* 2008;371:1527–37.
 31. Yang Y, Wu Z, Zhao T, Wang H, Zhao D, Zhang J, Wang Y, Ding Y, Qiu G. Adolescent Idiopathic Scoliosis and the Single-nucleotide Polymorphism of the Growth Hormone Receptor and IGF-1 Genes. *Orthopedics* 2009;32:411–6. [\[CrossRef\]](#)
 32. Sousa A, Fonseca I, Pichel F, Amaral TF. Effects of Posture and Body Mass Index on Body Girth Assessment. *Nutr Clin Pract* 2016;31:690–4. [\[CrossRef\]](#)